

Continuous Unitary Transformations

Vladimir. L. Safonov *

Center for Magnetic Recording Research, University of California - San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0401, U.S.A.

In the present time we observe a growing number of publications where the, so-called, flow equations are successfully used to diagonalize Hamiltonians by means of an appropriate unitary transformation. Here we discuss and compare the flow equations (FE) method (proposed in 1994) with the method of one step continuous unitary transformations (OS CUT) (proposed in 1982). It is shown that the FE method can be considered as a generalization of the OS CUT approach to the case of parameter dependent generator. The OS CUT method gives *linear* differential equations for the diagonalization procedure. In the FE method the system of differential equations is *nonlinear*. Finally we discuss the generalization of idea of continuous unitary transformations for the case of quantum equations of motion (Heisenberg picture and density matrix).

I. INTRODUCTION

Unitary transformations play an extremely important role in physics. For example, with the help of unitary transformation sometimes it is possible to simplify a problem of interacting quasiparticles by eliminating “inconvenient” interaction terms from initial Hamiltonian and construct corresponding effective interaction terms.

In the present time we observe a growing number of publications (see, [1], [2], [3], [4], [5], [6], [7], [8], [9], [10], [11], [12], [13], [14]), where the, so-called, flow equations (FE) method is successfully used to diagonalize Hamiltonians by means of an appropriate unitary transformation. This method of continuous unitary transformations looks very convenient and universal. The transformed Hamiltonian appears as a result of solving differential equations on some formal parameter.

It should be noted that quite the same idea and an analogous method (Safonov, 1982 [15], 1983 [16]) to construct nonlinear unitary transformation was published far before the original paper on FE method (Wegner, 1994 [1]) and just did not attract enough attention of researches. For convenience, we shall call the approach of Refs. [15], [16] as a one-step continuous unitary transformation (OS CUT) method.

The goal of the present paper is to discuss and compare FE and OS CUT methods. As it will be shown later, the FE method is a generalization of OS CUT approach. The only difference between the methods is that

the unitary transformation generator is assumed to be dependent on a formal parameter in the FE method and the generator does not depend on the parameter in the OS CUT method. In the CUT method we obtain *linear* differential equations. In the FE method we should solve a *nonlinear* system of differential equations. Below we shall demonstrate both methods by several examples.

It is interesting that the OS CUT method idea can be applied to the case of quantum equations of motion. Recently Mišta and Filip [23] have developed a method of non-perturbative solutions of nonlinear Heisenberg equations. We shall briefly discuss this approach at the end of this paper.

II. CONTINUOUS UNITARY TRANSFORMATION METHOD

A. One step continuous unitary transformation

Here we shall formulate the key idea of the OS CUT method [15], [16]. A general form of unitary transformation can be written as

$$H(\theta) = e^{\theta R} H e^{-\theta R} \quad (1)$$

where θ is a formal parameter, R is an anti-Hermitian operator ($R^\dagger = -R$), H and $H(\theta)$ are the initial and transformed Hamiltonians, respectively. The expression (1) is the solution of the equation

$$\frac{d}{d\theta} H(\theta) = [R, H(\theta)] \quad (2)$$

with the initial condition $H(0) = H$. In this form the Hamiltonian is “rotated in an operator space around R .”

In order to solve this equation we should write the most general form of $H(\theta)$ as the expansion in terms of operator combinations with unknown θ -dependent coefficients. The most general form of R as anti-Hermitian operator combinations also should be used.

After substituting these general (for $H(\theta)$ and R) expressions into (2), one can obtain a set of *linear* differential equations by comparing coefficients in analogous operator compositions in both sides. Solving these equations with the initial conditions, we obtain the transformed Hamiltonian $H(\theta)$. In order to eliminate ‘inconvenient’ terms one needs to put their coefficients (for example, for $\theta = 1$) equal to zero. This condition defines the choice of R .

Usually it is simple to find the anti-Hermitian operator, which eliminates given non-diagonal term. The OS CUT

*e-mail: vsafonov@ucsd.edu

method first has been applied for the spin Hamiltonian diagonalization [15], [16]. Then it was successfully used in physics of nuclear spin waves [17], magnon-impurity interactions [19], theory of superconductivity [19] and for eliminating three-boson interactions [20].

B. Flow equations for Hamiltonians

The flow equations method [1] begins with a general form of a unitary transformation:

$$H(\ell) = U(\ell) H U^\dagger(\ell), \quad (3)$$

where ℓ is a continuous flow parameter. The Hamiltonian $H(\ell)$ is transformed from the initial Hamiltonian $H(0) = H$. Differentiation (3) yields

$$\frac{d}{d\ell} H(\ell) = [\eta(\ell), H(\ell)] \quad (4)$$

with the generator

$$\eta(\ell) = \frac{dU(\ell)}{d\ell} U^\dagger(\ell) = -\eta^\dagger(\ell). \quad (5)$$

The generator $\eta(\ell)$ should be chosen in such a way, that the off-diagonal matrix elements decay. A good choice is defined by

$$\eta(\ell) = [H_d(\ell), H(\ell)], \quad (6)$$

where $H_d(\ell)$ is the diagonal part of $H(\ell)$ (comments ^{1,2}).

One can see that the FE method utilizes the same key tool (first proposed in the OS CUT method): the unitary transformation is represented as a solution of differential equations on a formal parameter. With an accuracy of notations the only difference between FE procedure and the OS CUT method is the parameter dependence of the generator $\eta(\ell)$. If η does not depend on ℓ we have $U(\ell) = \exp(\eta\ell)$. This case, obviously, exactly corresponds to the OS CUT method. If $U(\ell) = \exp(RF(\ell))$, where R is the parameter independent anti-Hermitian operator and $F(\ell)$ some function on ℓ , then $\eta(\ell) = (dF(\ell)/d\ell)R$. The Eq.(4) in this case can be written as

$$\frac{d}{(dF(\ell)/d\ell)d\ell} H(\ell) = [R, H(\ell)]. \quad (7)$$

This equation is reduced to Eq.(2) using a simple change of variable $d\theta = (dF(\ell)/d\ell)d\ell$. For $U(\ell) = \exp(R_1 F_1(\ell) + R_2 F_2(\ell))$ with commuting R_1 and R_2 anti-Hermitian operators one has $\eta(\ell) = (dF_1(\ell)/d\ell)R_1 +$

¹A rigorous criterion that this choice is most optimal seems has not been obtained.

²The condition of diagonality and the condition that $\eta(\ell) = [H_d(\ell), H(\ell)] = 0$ are not equivalent to each other.

$(dF_2(\ell)/d\ell)R_2$. The diagonalization procedure in this case can be considered as a two transformations with $U_1(\ell) = \exp(R_1 F_1(\ell))$ and $U_2(\ell) = \exp(R_2 F_2(\ell))$ unitary operators. Each transformation is described by (7) with $R_1, F_1(\ell)$, or $R_2, F_2(\ell)$.

In the most general case the ℓ -dependent generator transformation can not be reduced to the parameter independent case. In such a situation the FE method can give principally different results than the OS CUT method. Let us consider several examples.

III. QUADRATIC FORM DIAGONALIZATION

Stein [4] has considered the following quadratic Hamiltonian:

$$H_1(\ell) = \sum_{\mathbf{q}} \left(f_{\mathbf{q}}(\ell) (a_{\mathbf{q}}^\dagger a_{\mathbf{q}} + b_{\mathbf{q}}^\dagger b_{\mathbf{q}}) + g_{\mathbf{q}}(\ell) (a_{\mathbf{q}}^\dagger b_{-\mathbf{q}}^\dagger + a_{\mathbf{q}} b_{-\mathbf{q}}) \right), \quad (8)$$

where $a_{\mathbf{q}}^\dagger, a_{\mathbf{q}}$ and $b_{\mathbf{q}}^\dagger, b_{\mathbf{q}}$ are the creation and annihilation Bose operators. Applying flow equation

$$\frac{d}{d\ell} H_1(\ell) = [\eta_0(\ell), H_1(\ell)] \quad (9)$$

with

$$\eta_0(\ell) = \frac{1}{2} \sum_{\mathbf{q}} g_{\mathbf{q}}(\ell) (a_{\mathbf{q}}^\dagger b_{-\mathbf{q}}^\dagger - a_{\mathbf{q}} b_{-\mathbf{q}}), \quad (10)$$

the following *nonlinear* differential equations have been obtained:

$$\frac{d}{d\ell} f_{\mathbf{q}}(\ell) = -g_{\mathbf{q}}^2(\ell), \quad \frac{d}{d\ell} g_{\mathbf{q}}(\ell) = -f_{\mathbf{q}}(\ell) g_{\mathbf{q}}(\ell). \quad (11)$$

The solutions of (11) are

$$f_{\mathbf{q}}(\ell) = \epsilon_{\mathbf{q}} \coth(\ell \epsilon_{\mathbf{q}} + \ell_0(\mathbf{q})), \quad g_{\mathbf{q}}(\ell) = \frac{\epsilon_{\mathbf{q}} \operatorname{sgn}(\gamma_{\mathbf{q}})}{\sinh(\ell \epsilon_{\mathbf{q}} + \ell_0(\mathbf{q}))}, \quad (12)$$

where $\epsilon_{\mathbf{q}} = (1 - \gamma_{\mathbf{q}}^2)^{1/2}$, $\ell_0(\mathbf{q}) = \frac{1}{2} \ln \left(\frac{1 + \epsilon_{\mathbf{q}}}{1 - \epsilon_{\mathbf{q}}} \right)$ and

$$\epsilon_{\mathbf{q}} = \sqrt{f_{\mathbf{q}}^2(\ell) - g_{\mathbf{q}}^2(\ell)} = \sqrt{f_{\mathbf{q}}^2(0) - g_{\mathbf{q}}^2(0)} \quad (13)$$

is the spectrum of the diagonal (at $\ell \rightarrow \infty$) Hamiltonian (8).

Let us now consider the analogous diagonalization procedure in the framework of the OS CUT method. For convenience, the same notations will be used. Simple analysis shows that the only anti-Hermitian operator that can eliminate $a_{\mathbf{q}}^\dagger b_{-\mathbf{q}}^\dagger + a_{\mathbf{q}} b_{-\mathbf{q}}$ in (8) is $a_{\mathbf{q}}^\dagger b_{-\mathbf{q}}^\dagger - a_{\mathbf{q}} b_{-\mathbf{q}}$. Thus, the generator (independent on ℓ) should be taken in the form:

$$R = \frac{1}{2} \sum_{\mathbf{q}} G_{\mathbf{q}} (a_{\mathbf{q}}^{\dagger} b_{-\mathbf{q}}^{\dagger} - a_{\mathbf{q}} b_{-\mathbf{q}}), \quad (14)$$

where $G_{\mathbf{q}}$ is unknown function independent on ℓ . From $dH_1(\ell)/d\ell = [R, H_1(\ell)]$ it is simple to obtain *linear* differential equations

$$\frac{d}{d\ell} f_{\mathbf{q}}(\ell) = -G_{\mathbf{q}} g_{\mathbf{q}}(\ell), \quad \frac{d}{d\ell} g_{\mathbf{q}}(\ell) = -G_{\mathbf{q}} f_{\mathbf{q}}(\ell). \quad (15)$$

The solution of (15) is simple:

$$f_{\mathbf{q}}(\ell) = \frac{f_{\mathbf{q}}(0) + g_{\mathbf{q}}(0)}{2} e^{-G_{\mathbf{q}}\ell} + \frac{f_{\mathbf{q}}(0) - g_{\mathbf{q}}(0)}{2} e^{G_{\mathbf{q}}\ell}, \quad (16)$$

$$g_{\mathbf{q}}(\ell) = \frac{f_{\mathbf{q}}(0) + g_{\mathbf{q}}(0)}{2} e^{-G_{\mathbf{q}}\ell} - \frac{f_{\mathbf{q}}(0) - g_{\mathbf{q}}(0)}{2} e^{G_{\mathbf{q}}\ell}.$$

Let the Hamiltonian (8) be diagonal at $\ell = 1$: $g_{\mathbf{q}}(1) = 0$. From this condition we immediately find the unknown function $G_{\mathbf{q}}$:

$$G_{\mathbf{q}} = \frac{1}{2} \ln \left(\frac{f_{\mathbf{q}}(0) + g_{\mathbf{q}}(0)}{f_{\mathbf{q}}(0) - g_{\mathbf{q}}(0)} \right) \quad (17)$$

and the spectrum

$$\epsilon_{\mathbf{q}} = f_{\mathbf{q}}(1) = \sqrt{f_{\mathbf{q}}^2(0) - g_{\mathbf{q}}^2(0)}. \quad (18)$$

Thus, we see that in the case of exact diagonalization of quadratic Hamiltonian the results of FE and OS CUT methods coincide with each other.

IV. ELECTRON-PHONON COUPLING

Consider now the elimination of electron-phonon interaction to obtain the effective electron-electron scattering in the framework of OS CUT and FE methods. The Hamiltonian is (see, [3]):

$$H = H_0 + H_{e-ph} + H_{e-e}, \quad (19)$$

where

$$H_0 = \sum_{\mathbf{q}} \omega_{\mathbf{q}} a_{\mathbf{q}}^{\dagger} a_{\mathbf{q}} + \sum_{\mathbf{k}} \varepsilon_{\mathbf{k}} c_{\mathbf{k}}^{\dagger} c_{\mathbf{k}} \quad (20)$$

describes free phonons ($a_{\mathbf{q}}^{\dagger}$, $a_{\mathbf{q}}$) and electrons ($c_{\mathbf{k}}^{\dagger}$, $c_{\mathbf{k}}$),

$$H_{e-ph}(\ell) = \sum_{\mathbf{k}, \mathbf{q}} M_{\mathbf{k}, \mathbf{q}}(\ell) a_{-\mathbf{q}}^{\dagger} c_{\mathbf{k}+\mathbf{q}}^{\dagger} c_{\mathbf{k}} + \text{h.c.} \quad (21)$$

describes the electron-phonon interaction and

$$H_{e-e}(\ell) = \sum_{\mathbf{k}, \mathbf{k}', \mathbf{q}} V_{\mathbf{k}, \mathbf{k}', \mathbf{q}}(\ell) c_{\mathbf{k}+\mathbf{q}}^{\dagger} c_{\mathbf{k}'-\mathbf{q}}^{\dagger} c_{\mathbf{k}'} c_{\mathbf{k}} \quad (22)$$

is the Hamiltonian of electron-electron interaction. The initial conditions are:

$$M_{\mathbf{k}, \mathbf{q}}(\ell = 0) = M_{\mathbf{q}}, \quad V_{\mathbf{k}, \mathbf{k}', \mathbf{q}}(\ell = 0) = V_{\mathbf{k}, \mathbf{k}', \mathbf{q}}(0). \quad (23)$$

Let us first consider the diagonalization (elimination of (21)) in the framework of OS CUT method. From a simple analysis it follows that the parameter independent generator should be taken in the form:

$$R = \sum_{\mathbf{k}, \mathbf{q}} R_{\mathbf{k}, \mathbf{q}} a_{-\mathbf{q}}^{\dagger} c_{\mathbf{k}+\mathbf{q}}^{\dagger} c_{\mathbf{k}} - \text{h.c.} \quad (24)$$

The diagonalization procedure gives the following *linear* differential equations:

$$\frac{\partial M_{\mathbf{k}, \mathbf{q}}(\ell)}{\partial \ell} = -\alpha_{\mathbf{k}, \mathbf{q}} R_{\mathbf{k}, \mathbf{q}} \quad (25)$$

and for the electron pairs (with zero momentum)

$$\frac{\partial V_{\mathbf{k}, -\mathbf{k}, \mathbf{q}}(\ell)}{\partial \ell} = -R_{\mathbf{k}, \mathbf{q}} M_{-\mathbf{k}-\mathbf{q}, \mathbf{q}}(\ell) - R_{-\mathbf{k}-\mathbf{q}, \mathbf{q}} M_{\mathbf{k}, \mathbf{q}}(\ell). \quad (26)$$

From (25) we obtain

$$M_{\mathbf{k}, \mathbf{q}}(\ell) = M_{\mathbf{k}, \mathbf{q}}(0) - \alpha_{\mathbf{k}, \mathbf{q}} R_{\mathbf{k}, \mathbf{q}} \ell. \quad (27)$$

Let the electron-phonon Hamiltonian (21) vanishes at $\ell = 1$: $M_{\mathbf{k}, \mathbf{q}}(1) = 0$. From this condition we immediately find

$$R_{\mathbf{k}, \mathbf{q}} = M_{\mathbf{k}, \mathbf{q}}(0) / \alpha_{\mathbf{k}, \mathbf{q}}. \quad (28)$$

Simple solution of (26) with (27), (28) and initial conditions (23) gives

$$V_{\mathbf{k}, -\mathbf{k}, \mathbf{q}}(1) = V_{\mathbf{k}, -\mathbf{k}, \mathbf{q}}(0) - \frac{M_{\mathbf{q}}^2 \omega_{\mathbf{q}}}{\omega_{\mathbf{q}}^2 - (\varepsilon_{\mathbf{k}+\mathbf{q}} - \varepsilon_{\mathbf{k}})^2}. \quad (29)$$

This result corresponds to the Fröhlich's effective interaction obtained in the second order of the perturbation theory (see, e.g., [21]).

Let us now consider the FE method. The generator is taken in the form [3]:

$$\eta(\ell) = [H_0, H_{e-ph}(\ell)] \quad (30)$$

$$= \sum_{\mathbf{k}, \mathbf{q}} M_{\mathbf{k}, \mathbf{q}}(\ell) \alpha_{\mathbf{k}, \mathbf{q}} a_{-\mathbf{q}}^{\dagger} c_{\mathbf{k}+\mathbf{q}}^{\dagger} c_{\mathbf{k}} - \text{h.c.}$$

with the energy difference

$$\alpha_{\mathbf{k}, \mathbf{q}} = \varepsilon_{\mathbf{k}+\mathbf{q}} - \varepsilon_{\mathbf{k}} + \omega_{\mathbf{q}}. \quad (31)$$

This generator yields several contributions to $dH(\ell)/d\ell = [\eta(\ell), H(\ell)]$. The contribution to the change of $M_{\mathbf{k}, \mathbf{q}}(\ell)$ results from $[\eta(\ell), H_0]$

$$\frac{\partial M_{\mathbf{k}, \mathbf{q}}(\ell)}{\partial \ell} = -\alpha_{\mathbf{k}, \mathbf{q}}^2 M_{\mathbf{k}, \mathbf{q}}(\ell) \quad (32)$$

with the solution

$$M_{\mathbf{k}, \mathbf{q}}(\ell) = M_{\mathbf{q}} \exp(-\alpha_{\mathbf{k}, \mathbf{q}}^2 \ell), \quad (33)$$

where $M_{\mathbf{q}}$ is the initial electron-phonon coupling. For $\alpha_{\mathbf{k},\mathbf{q}} \neq 0$ the solution (33) eliminates the electron-phonon interaction from the Hamiltonian at $\ell \rightarrow \infty$. The contribution to the effective electron-electron interaction is obtained from $[\eta(\ell), H_{e-ph}(\ell)]$. For the electron pairs (with zero momentum) we have the following *nonlinear* differential equation:

$$\frac{\partial V_{\mathbf{k},-\mathbf{k},\mathbf{q}}(\ell)}{\partial \ell} = -(\alpha_{\mathbf{k},\mathbf{q}} + \alpha_{-\mathbf{k}-\mathbf{q},\mathbf{q}})M_{\mathbf{k},\mathbf{q}}(\ell)M_{-\mathbf{k}-\mathbf{q},\mathbf{q}}(\ell) \quad (34)$$

with the solution

$$V_{\mathbf{k},-\mathbf{k},\mathbf{q}}(\infty) = V_{\mathbf{k},-\mathbf{k},\mathbf{q}}(0) - \frac{M_{\mathbf{q}}^2 \omega_{\mathbf{q}}}{\omega_{\mathbf{q}}^2 + (\varepsilon_{\mathbf{k}+\mathbf{q}} - \varepsilon_{\mathbf{k}})^2}. \quad (35)$$

This result differs from (29) (plus sign between the two squares in the denominator). The difference is explained as: “perturbation theory for Hamiltonians is not uniquely defined” [3].

In the FE method a choice of $\eta(\ell)$ is motivated to be defined by $[H_0, H_{e-ph}(\ell)]$ (30). However, a rigorous criterion that this choice (or, another one) is most optimal for a given problem seems has not yet been formulated.

V. THREE-BOSON HAMILTONIAN

In this example we consider the elimination of forbidden three-boson terms from the Hamiltonian

$$H(\ell) = \sum_{\mathbf{k}} \varepsilon_{\mathbf{k}} b_{\mathbf{k}}^{\dagger} b_{\mathbf{k}} + H_3(\ell) + H_4(\ell), \quad (36)$$

where

$$H_3(\ell) = \frac{1}{3} \sum_{\mathbf{k},\mathbf{q}} \{ \Psi_{\mathbf{k},\mathbf{q}}(\ell) b_{\mathbf{k}}^{\dagger} b_{\mathbf{q}}^{\dagger} b_{-\mathbf{k}-\mathbf{q}}^{\dagger} + \text{h.c.} \} \quad (37)$$

and

$$H_4(\ell) = \frac{1}{2} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} \Phi_{\mathbf{k}_1, \mathbf{k}_2; \mathbf{k}_3, \mathbf{k}_4}(\ell) b_{\mathbf{k}_1}^{\dagger} b_{\mathbf{k}_2}^{\dagger} b_{\mathbf{k}_3} b_{\mathbf{k}_4} \times \Delta(\mathbf{k}_1 + \mathbf{k}_2 - \mathbf{k}_3 - \mathbf{k}_4). \quad (38)$$

Here ℓ is a formal parameter and $\Delta(\mathbf{k})$ is the Kronecker delta function. For simplicity, we shall neglect the energy $\varepsilon_{\mathbf{k}}$ dependence on ℓ .

Using OS CUT method, we write the parameter-independent operator in the form [20]:

$$R = \sum_{\mathbf{k},\mathbf{q}} (R_{\mathbf{k},\mathbf{q}} b_{\mathbf{k}}^{\dagger} b_{\mathbf{q}}^{\dagger} b_{-\mathbf{k}-\mathbf{q}}^{\dagger} - \text{h.c.}). \quad (39)$$

Calculating the commutators $[R, \sum_{\mathbf{k}} \varepsilon_{\mathbf{k}} b_{\mathbf{k}}^{\dagger} b_{\mathbf{k}}]$ and $[R, H_3(\ell)]$, we obtain the following linear differential equations:

$$\frac{d}{d\ell} \Psi_{\mathbf{k},\mathbf{q}}(\ell) = -3(\varepsilon_{\mathbf{k}} + \varepsilon_{\mathbf{q}} + \varepsilon_{-\mathbf{k}-\mathbf{q}}) R_{\mathbf{k},\mathbf{q}}, \quad (40)$$

$$\begin{aligned} \frac{d}{d\ell} \Phi_{\mathbf{k}_1, \mathbf{k}_2; \mathbf{k}_3, \mathbf{k}_4}(\ell) = & -6R_{\mathbf{k}_1, \mathbf{k}_2} \Psi_{\mathbf{k}_3, \mathbf{k}_4}^*(\ell) \\ & -6R_{\mathbf{k}_3, \mathbf{k}_4}^* \Psi_{\mathbf{k}_1, \mathbf{k}_2}(\ell). \end{aligned} \quad (41)$$

Solving these equations with the initial conditions $\Psi_{\mathbf{k}_1, \mathbf{k}_2}(\ell = 0) = \Psi_{\mathbf{k}_1, \mathbf{k}_2}$, $\Phi_{\mathbf{k}_1, \mathbf{k}_2; \mathbf{k}_3, \mathbf{k}_4}(\ell = 0) = 0$, and demanding $\Psi_{\mathbf{k}_1, \mathbf{k}_2}(\ell = 1) = 0$, we obtain

$$R_{\mathbf{k},\mathbf{q}} = \frac{\Psi_{\mathbf{k},\mathbf{q}}}{3(\varepsilon_{\mathbf{k}} + \varepsilon_{\mathbf{q}} + \varepsilon_{-\mathbf{k}-\mathbf{q}})} \quad (42)$$

and

$$\begin{aligned} \Phi_{\mathbf{k}_1, \mathbf{k}_2; \mathbf{k}_3, \mathbf{k}_4}(1) = & -\Psi_{\mathbf{k}_1, \mathbf{k}_2} \Psi_{\mathbf{k}_3, \mathbf{k}_4}^* \left(\frac{1}{\varepsilon_{\mathbf{k}_1} + \varepsilon_{\mathbf{k}_2} + \varepsilon_{-\mathbf{k}_1-\mathbf{k}_2}} \right. \\ & \left. + \frac{1}{\varepsilon_{\mathbf{k}_3} + \varepsilon_{\mathbf{k}_4} + \varepsilon_{-\mathbf{k}_3-\mathbf{k}_4}} \right). \end{aligned} \quad (43)$$

It should be noted that this result for the effective four-boson interaction coincides (with the additional condition $\Psi_1 = \Psi_1^*$) with the result of classical diagonalization of (37) (when $b_{\mathbf{k}}^{\dagger}$ and $b_{\mathbf{k}}$ are *c*-numbers) by canonical quasi-linear transformation [22].

Let us now consider the same problem in the framework of FE method. The (parameter-dependent) generator in this case is:

$$\eta(\ell) = \frac{1}{3} \sum_{\mathbf{k},\mathbf{q}} \{ \beta_{\mathbf{k},\mathbf{q}} \Psi_{\mathbf{k},\mathbf{q}}(\ell) b_{\mathbf{k}}^{\dagger} b_{\mathbf{q}}^{\dagger} b_{-\mathbf{k}-\mathbf{q}}^{\dagger} - \text{h.c.} \}, \quad (44)$$

where

$$\beta_{\mathbf{k},\mathbf{q}} = \varepsilon_{\mathbf{k}} + \varepsilon_{\mathbf{q}} + \varepsilon_{-\mathbf{k}-\mathbf{q}}. \quad (45)$$

Corresponding differential equation for the three-boson amplitude has the form:

$$\frac{d}{d\ell} \Psi_{\mathbf{k},\mathbf{q}}(\ell) = -\beta_{\mathbf{k},\mathbf{q}}^2 \Psi_{\mathbf{k},\mathbf{q}}(\ell) \quad (46)$$

with the solution (compare with (33)):

$$\Psi_{\mathbf{k},\mathbf{q}}(\ell) = \Psi_{\mathbf{k},\mathbf{q}} \exp(-\beta_{\mathbf{k},\mathbf{q}}^2 \ell). \quad (47)$$

The differential equation for the four-boson effective amplitude in this case is

$$\frac{d\Phi_{\mathbf{k}_1, \mathbf{k}_2; \mathbf{k}_3, \mathbf{k}_4}(\ell)}{d\ell} = -2(\beta_{\mathbf{k}_1, \mathbf{k}_2} + \beta_{\mathbf{k}_3, \mathbf{k}_4}) \Psi_{\mathbf{k}_1, \mathbf{k}_2}(\ell) \Psi_{\mathbf{k}_3, \mathbf{k}_4}^*(\ell). \quad (48)$$

The solution at $\ell \rightarrow \infty$ (when $\Psi_{\mathbf{k},\mathbf{q}}(\ell)$ vanishes) has the form

$$\Phi_{\mathbf{k}_1, \mathbf{k}_2; \mathbf{k}_3, \mathbf{k}_4}(1) = -2\Psi_{\mathbf{k}_1, \mathbf{k}_2} \Psi_{\mathbf{k}_3, \mathbf{k}_4}^* \frac{\beta_{\mathbf{k}_1, \mathbf{k}_2} + \beta_{\mathbf{k}_3, \mathbf{k}_4}}{\beta_{\mathbf{k}_1, \mathbf{k}_2}^2 + \beta_{\mathbf{k}_3, \mathbf{k}_4}^2}. \quad (49)$$

We see that this formula differs from the OS CUT result (43) and therefore (49) does not correspond to the classical case [22].

VI. NON-PERTURBATIVE SOLUTION OF QUANTUM EQUATIONS

A. Heisenberg picture

The Heisenberg equation for an operator $A(t)$

$$i\hbar \frac{d}{dt} A(t) = [A(t), H(t)] \quad (50)$$

can be rewritten in the form:

$$\frac{d}{dt} A(t) = [iH(t)/\hbar, A(t)]. \quad (51)$$

Formally this equation is similar to (2) (or, (4) for time-dependent Hamiltonian): time t plays a role of a formal parameter and $iH(t)/\hbar$ is an anti-Hermitian operator. As in the CUT method, we can write the most general form of $A(t)$ as the expansion in terms of operator combinations with unknown time-dependent coefficients. After substituting these form into (51) with a given $iH(t)/\hbar$, one can obtain a set of differential equations by comparing coefficients in analogous operator compositions in both sides. These equations will be linear for time-independent H and nonlinear for time-dependent $H(t)$. Then we solve these equations with the initial conditions defined by $A(t) = A(0)$. Recently Mišta and Filip [23] have developed this idea as a method of non-perturbative solutions of nonlinear Heisenberg equations with an illustrative example of two coupled harmonic oscillators.

B. Density matrix

It is obvious that a quite analogous idea can be applied for the density matrix. In this case we can write the density matrix equation as

$$\frac{d}{dt} \rho(t) = [-iH(t)/\hbar, \rho(t)], \quad (52)$$

where $-iH(t)/\hbar$ plays a role of “transformation generator”.

As an illustration consider a system of nuclear spins $\mathbf{I}_j$ with a Hamiltonian $H = H_0 + H_{int}$, where $H_0 = \mu B \sum_j I_j^z$ is the Zeeman energy and H_{int} describes spin-spin interactions. As an initial state assume the transverse magnetization. In the high temperature approximation one can write $\rho(0) = 1 - \alpha \sum_j I_j^x$. Then the density matrix can be represented as $\rho(t) = 1 - \sum_{n=1}^{\infty} \Delta \rho_n(t)$, where $\Delta \rho_n(t)$ describes compositions of n spins with time dependent coefficients, e.g., $\Delta \rho_1(t) = \sum_j (X_j(t) I_j^x + Y_j(t) I_j^y)$. From $[H_0, -\Delta \rho_1(t)]$ we obtain differential equations $dX_j(t)/dt = -\omega_0 Y_j(t)$, $dY_j(t)/dt = \omega_0 X_j(t)$ and solutions $X_j(t) = \alpha \cos \omega_0 t$, $Y_j(t) = \alpha \sin \omega_0 t$, where $\omega_0 = \mu B/\hbar$ is the nuclear magnetic resonance frequency. Contributions to two spin motions can be found from $[H_0, -\Delta \rho_2(t)] + [H_{int}, -\Delta \rho_1(t)]$ and so on.

This example can demonstrate how the order stored in magnetization is transformed to higher-order spin correlations.

The author wish to thank G. S. Uhrig and A. K. Khitrin for valuable comments.

-
- [1] F. Wegner, Ann. Physik **3**, 77 (1994).
 - [2] S. K. Kehrein, A. Mielke, and P. Neu, Z. Phys. B **99**, 269 (1996).
 - [3] F. Wegner, Philosophical Magazine B **77**, 1249 (1998)
 - [4] J. Stein, Eur. Phys. J. B **5**, 193 (1998).
 - [5] C. P. Moca, I. Tifrea, and M. Crisan, J. of Superconductivity **12**, 399 (1999).
 - [6] J. Stein, Europhys. Lett. **50**, 68 (2000).
 - [7] E. L. Gubankova, C.-R. Ji, and S. R. Cotanch, Phys. Rev. D **62**, 074001 (2000).
 - [8] W. Hofstetter and S. Kehrein, Phys. Rev. B **63**, 140402(R) (2001).
 - [9] C. Knetter, K. P. Schmidt, M. Grüninger, and G. S. Uhrig, Phys. Rev. Lett. **87**, 167204 (2001).
 - [10] T. Domański, Eur. Phys. J. B **23**, 49 (2001).
 - [11] C. Raas, A. Bühler, and G. S. Uhrig, Eur. Phys. J. B **21**, 369 (2001).
 - [12] Y. Ohira and K. Imafuku, quant-ph/ 0201005.
 - [13] S. R. White, cond-mat/0201346.
 - [14] T. Domański and J. Ranninger, cond-mat/ 0202010.
 - [15] V. L. Safonov, Preprint KIAE - 3691/1, Moscow, August 8, 1982 (in Russian).
 - [16] V. L. Safonov, Phys. Lett. A **97**, 164 (1983).
 - [17] A. V. Andrienko, V. I. Ozhogin, V. L. Safonov, and A. Yu. Yakubovskii, Zh. Eksp. Teor. Fiz. **84**, 1158 (1983) [Sov. Phys. JETP **57**, 673 (1983)].
 - [18] V. L. Safonov and R. M. Farzetdinova, J. Magn. Magn. Mater. **98**, L235 (1991).
 - [19] V. L. Safonov, Phys. Stat. Solidi (b) **174**, 223 (1992).
 - [20] Q. Shi, V. L. Safonov, M. Mino, and H. Yamazaki, Phys. Lett. A **238**, 258 (1998).
 - [21] C. Kittel, *Quantum Theory of Solids* (Wiley, New York, 1963), p.152.
 - [22] V. P. Krasitskii, Sov. Phys.-JETP **71**, 921 (1990).
 - [23] L. Mišta, Jr. and R. Filip, J. Phys. A: Math. Gen. **34**, 5603 (2001).